

Planck 2015 results: XXVII. The second Planck catalogue of Sunyaev-Zeldovich sources

Ade P., Aghanim N., Arnaud M., Ashdown M., Aumont J., Baccigalupi C., Banday A., Barreiro R., Barrena R., Bartlett J., Bartolo N., Battaner E., Battye R., Benabed K., Benoît A., Benoit-Lévy A., Bernard J., Bersanelli M., Bielewicz P., Bikmaev I., Böhringer H., Bonaldi A., Bonavera L., Bond J., Borrill J., Bouchet F., Bucher M., Burenin R., Burigana C., Butler R., Calabrese E., Cardoso J., Carvalho P., Catalano A., Challinor A., Chamballu A., Chary R., Chiang H., Chon G., Christensen P., Clements D., Colombi S., Colombo L., Combet C., Comis B., Couchot F., Coulais A., Crill B., Curto A., Cuttaia F., Dahle H., Danese L., Davies R., Davis R., De Bernardis P., De Rosa A., De Zotti G., Delabrouille J., Désert F., Dickinson C., Diego J., Dolag K., Dole H., Donzelli S.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2016 ESO. We present the all-sky Planck catalogue of Sunyaev-Zeldovich (SZ) sources detected from the 29 month full-mission data. The catalogue (PSZ2) is the largest SZ-selected sample of galaxy clusters yet produced and the deepest systematic all-sky survey of galaxy clusters. It contains 1653 detections, of which 1203 are confirmed clusters with identified counterparts in external data sets, and is the first SZ-selected cluster survey containing >103 confirmed clusters. We present a detailed analysis of the survey selection function in terms of its completeness and statistical reliability, placing a lower limit of 83% on the purity. Using simulations, we find that the estimates of the SZ strength parameter Y_{5R500} are robust to pressure-profile variation and beam systematics, but accurate conversion to Y_{500} requires the use of prior information on the cluster extent. We describe the multi-wavelength search for counterparts in ancillary data, which makes use of radio, microwave, infra-red, optical, and X-ray data sets, and which places emphasis on the robustness of the counterpart match. We discuss the physical properties of the new sample and identify a population of low-redshift X-ray under-luminous clusters revealed by SZ selection. These objects appear in optical and SZ surveys with consistent properties for their mass, but are almost absent from ROSAT X-ray selected samples.

<http://dx.doi.org/10.1051/0004-6361/201525823>

Keywords

Catalogs, Cosmology: observations, Galaxies: clusters: general